

Correcting for the alias effect when measuring the power spectrum using FFT

Y.P. Jing

*Shanghai Astronomical Observatory, the Partner Group of MPI für Astrophysik,
Nandan Road 80, Shanghai 200030, China*

ABSTRACT

Because of mass assignment onto grid points in the measurement of the power spectrum using the Fast Fourier Transform (FFT), the raw power spectrum $\langle |\delta^f(k)|^2 \rangle$ estimated with FFT is not the same as the true power spectrum $P(k)$. In this paper, we derive the formula which relates $\langle |\delta^f(k)|^2 \rangle$ to $P(k)$. For a sample of N discrete objects, the formula reads: $\langle |\delta^f(k)|^2 \rangle = \sum_{\mathbf{n}} [|W(\mathbf{k} + 2k_N \mathbf{n})|^2 P(\mathbf{k} + 2k_N \mathbf{n}) + 1/N |W(\mathbf{k} + 2k_N \mathbf{n})|^2]$, where $W(\mathbf{k})$ is the Fourier transform of the mass assignment function $W(\mathbf{r})$, k_N is the Nyquist wavenumber, and $\mathbf{n}$ is an integer vector. The formula is different from that in some of previous works where the summation over $\mathbf{n}$ is neglected. For the NGP, CIC and TSC assignment functions, we show that the shot noise term $\sum_{\mathbf{n}} 1/N |W(\mathbf{k} + 2k_N \mathbf{n})|^2$ can be expressed by simple analytical functions. To reconstruct $P(k)$ from the alias sum $\sum_{\mathbf{n}} |W(\mathbf{k} + 2k_N \mathbf{n})|^2 P(\mathbf{k} + 2k_N \mathbf{n})$, we propose an iterative method. We test the method by applying it to an N-body simulation sample, and show that the method can successfully recover $P(k)$. The discussion is further generalized to samples with observational selection effects.

Subject headings: methods: data analysis - methods: statistical - galaxies: clustering - large-scale structure of Universe

1. INTRODUCTION

The power spectrum $P(k)$ of the spatial cosmic density distribution is an important quantity in galaxy formation theories. $P(k)$ on large scales is a direct measure of the primordial density fluctuation, and $P(k)$ on small scales carries information of later non-linear evolution, therefore measuring $P(k)$ can serve to distinguish between different theoretical models.

The power spectrum $P(k)$ as a clustering measure has already been applied by many authors to observational samples of galaxies and of clusters of galaxies, including the CfA and Perseus-Pisces redshift surveys (Baumgart & Fry 1991), the radio galaxy survey (Peacock & Nicholson 1991), the IRAS QDOT survey (Kaiser 1991), the galaxy distribution in the nearby superclusters (Gramann & Einasto 1992), the SSRS redshift sample (Park, Gott, & da Costa 1992), the CfA and SSRS extensions (Vogeley et al. 1992; da Costa et al. 1994), the 2Jy IRAS survey (Jing & Valdarnini 1993), the 1.2Jy IRAS survey (Fisher et al. 1993) and the redshift samples of Abell clusters (Jing & Valdarnini 1993; Peacock & West 1992). The power spectrum is also widely measured for cosmological N-body simulations, since it can easily characterize the linear and non-linear evolutions of the density perturbation (e.g. Davis et al. 1985).

All of these workers except Fisher et al., have used the Fast Fourier Transformation (FFT) technique to make the Fourier Transforms (FT). In fact, one can use the direct summation [Eq. (5) below] to measure the power spectrum for a sample of a few thousand objects (Fisher et al. 1993). However it appears to be more convenient for most workers to use FFT packages (now available on many computers) to analyze the power spectrum. This is probably the reason why most of the previous statistical studies of the power spectrum have used FFT. For N-body simulations, one has to use FFT to obtain the power spectrum, since normally there are more than a million particles. When using FFT, one needs to collect density values $\rho(\mathbf{r}_g)$ on a grid from a density field $\rho(\mathbf{r})$ (or a particle distribution), which is usually called ‘mass assignment’. The mass assignment is equivalent to convolving the density field by a given assignment function $W(\mathbf{r})$ and sampling the convolved density field on a finite number of grid points. The FT of $\rho(\mathbf{r}_g)$ generally is not equal to the FT of $\rho(\mathbf{r})$, and the power spectrum estimated directly from the FT of $\rho(\mathbf{r}_g)$ is a biased one. The smoothing effect has already been considered in many previous works (e.g. Baumgart & Fry 1991; Jing & Valdarnini 1993; Scoccimarro et al. 1998), but the sampling effect has not.

In this paper, we derive the formulae which express these effects of the mass assignment on the estimated power spectrum. A procedure is proposed to correct for these effects, in order to recover the true power spectrum. The procedure is tested and shown to be very successful with a simulation sample.

2. Formulae

Let us first recall the definition of the power spectrum $P(k)$. Let $\rho(\mathbf{r})$ be the cosmic density field and $\bar{\rho}$ the mean density. The density field can be expressed with a dimensionless

field $\delta(\mathbf{r})$ (which is usually called the density contrast):

$$\delta(\mathbf{r}) = \frac{\rho(\mathbf{r}) - \bar{\rho}}{\bar{\rho}}. \quad (1)$$

Based on the cosmological principle, one can imagine that $\rho(\mathbf{r})$ is periodic in some large rectangular volume V_μ . The FT of $\delta(\mathbf{r})$ is then defined as:

$$\delta(\mathbf{k}) = \frac{1}{V_\mu} \int_{V_\mu} \delta(\mathbf{r}) e^{i\mathbf{r} \cdot \mathbf{k}} d\mathbf{r}, \quad (2)$$

and the power spectrum $P(k)$ is simply related to $\delta(\mathbf{k})$ by

$$P(k) \equiv \langle |\delta(\mathbf{k})|^2 \rangle, \quad (3)$$

where $\langle \dots \rangle$ means the ensemble average.

In practical measurement of $P(k)$ either for an extragalactic catalogue or for simulation data, the continuous density field $\rho(\mathbf{r})$ is sampled by only a finite number N of objects. In these cases, one needs to deal with the ‘discreteness’ effect arising from the Poisson shot noise. To show this, let’s consider an ideal case where no selection effect has been introduced into the sample (In fact, some kind of selection effects must exist in extragalactic catalogues. We will discuss it in Section 3). The number density distribution of objects can be expressed as $n(\mathbf{r}) = \sum_j \delta^D(\mathbf{r} - \mathbf{r}_j)$, where $\mathbf{r}_j$ is the coordinate of object j and $\delta^D(\mathbf{r})$ is the Dirac- δ function. In analogy with the continuous case, the FT of $n(\mathbf{r})$ is defined as:

$$\delta^d(\mathbf{k}) = \frac{1}{V_\mu \bar{n}} \int_{V_\mu} n(\mathbf{r}) e^{i\mathbf{r} \cdot \mathbf{k}} d\mathbf{r} - \delta_{\mathbf{k}, \mathbf{0}}^K, \quad (4)$$

where $\bar{n}$ is the global mean number density, the superscript d represents the *discrete* case of $\rho(\mathbf{r})$, and δ^K is the Kronecker delta. Following Peebles (1980, sections 36–41), we divide the volume V_μ into infinitesimal elements $\{dV_i\}$ with n_i objects inside dV_i . Then the above equation can be written as:

$$\delta^d(\mathbf{k}) = \frac{1}{N} \sum_i n_i e^{i\mathbf{r}_i \cdot \mathbf{k}} - \delta_{\mathbf{k}, \mathbf{0}}^K, \quad (5)$$

where N is $\bar{n}V_\mu$, the number of objects in V_μ . Since dV_i is taken so small that n_i is either 0 or 1, we have $n_i = n_i^2 = n_i^3 = \dots$, $\langle n_i \rangle = \bar{n}dV_i$ and $\langle n_i n_j \rangle_{i \neq j} = \bar{n}^2 dV_i dV_j [1 + \langle \delta(\mathbf{r}_i) \delta(\mathbf{r}_j) \rangle]$. We can find the ensemble average of $\delta^d(\mathbf{k}_1) \delta^{d*}(\mathbf{k}_2)$:

$$\langle \delta^d(\mathbf{k}_1) \delta^{d*}(\mathbf{k}_2) \rangle = \frac{1}{N^2} \sum_{i,j} \langle n_i n_j \rangle e^{i\mathbf{r}_i \cdot \mathbf{k}_1 - i\mathbf{r}_j \cdot \mathbf{k}_2} - \delta_{\mathbf{k}_1, \mathbf{0}}^K \delta_{\mathbf{k}_2, \mathbf{0}}^K$$

$$\begin{aligned}
&= \frac{1}{N^2} \sum_{i \neq j} \langle n_i n_j \rangle e^{i\mathbf{r}_i \cdot \mathbf{k}_1 - i\mathbf{r}_j \cdot \mathbf{k}_2} + \frac{1}{N^2} \sum_i \langle n_i \rangle e^{i\mathbf{r}_i \cdot (\mathbf{k}_1 - \mathbf{k}_2)} - \delta_{\mathbf{k}_1, \mathbf{0}}^K \delta_{\mathbf{k}_2, \mathbf{0}}^K \\
&= \langle \delta(\mathbf{k}_1) \delta^*(\mathbf{k}_2) \rangle + \frac{1}{N} \delta_{\mathbf{k}_1, \mathbf{k}_2}^K .
\end{aligned} \tag{6}$$

The last equation assumes $\mathbf{k}_1 \neq 0$ or $\mathbf{k}_2 \neq 0$. The true power spectrum is then:

$$P(k) \equiv \langle |\delta(\mathbf{k})|^2 \rangle = \langle |\delta^d(\mathbf{k})|^2 \rangle - \frac{1}{N} . \tag{7}$$

So the discreteness (or shot noise) effect is to introduce an additional term $1/N$ to the power spectrum $\langle |\delta^d(\mathbf{k})|^2 \rangle$. This fact is already well known to cosmologists. We present the above derivation because this method is useful in the following derivations.

In principle one can use the direct summation of Eq. (5) to measure the power spectrum for a sample of discrete objects. However, as described in Section 1, most of the previous statistical studies of the power spectrum have used FFT. Moreover it would be impossible to use direct summation to measure the power spectrum for an N-body simulation. The quantity computed by the FFT is

$$\delta^f(\mathbf{k}) = \frac{1}{N} \sum_{\mathbf{g}} n^f(\mathbf{r}_g) e^{i\mathbf{r}_g \cdot \mathbf{k}} - \delta_{\mathbf{k}, \mathbf{0}}^K , \tag{8}$$

where the superscript f denotes quantities in FFT. $n^f(\mathbf{r}_g)$ is the convolved density value on the $\mathbf{g}$ -th grid point $\mathbf{r}_g = \mathbf{g}H$ ($\mathbf{g}$ is an integer vector; H is the grid spacing):

$$n^f(\mathbf{r}_g) = \int n(\mathbf{r}) W(\mathbf{r} - \mathbf{r}_g) d\mathbf{r} , \tag{9}$$

where $W(\mathbf{r})$ is the mass assignment function.

Following Hockney and Eastwood (1981), Eq.(8) can be expressed in a more compact way by using the so-called ‘Sampling Function’. The sampling function $\Pi(\mathbf{r})$ is defined as a sum of the Dirac- δ functions spaced at unit length in all three spatial directions, i.e. $\Pi(\mathbf{r}) = \sum_{\{\mathbf{n}\}} \delta^D(\mathbf{r} - \mathbf{n})$, where $\mathbf{n}$ is an integer vector. Defining

$$n'^f(\mathbf{r}) \equiv \Pi\left(\frac{\mathbf{r}}{H}\right) \int n(\mathbf{r}_1) W(\mathbf{r}_1 - \mathbf{r}) d\mathbf{r}_1 \tag{10}$$

and constructing

$$\delta'^f(\mathbf{k}) = \frac{1}{N} \int n'^f(\mathbf{r}) e^{i\mathbf{r} \cdot \mathbf{k}} d\mathbf{r} - \delta_{\mathbf{k}, \mathbf{0}}^K , \tag{11}$$

one can easily prove

$$\delta'^f(\mathbf{k}) = \delta^f(\mathbf{k}) . \tag{12}$$

Therefore one can express $\delta^f(\mathbf{k})$ as [cf. Eqs. (10-12)]:

$$\delta^f(\mathbf{k}) = \frac{1}{N} \int_{V_\mu} \Pi\left(\frac{\mathbf{r}}{H}\right) \sum_i n_i W(\mathbf{r}_i - \mathbf{r}) e^{i\mathbf{r} \cdot \mathbf{k}} d\mathbf{r} - \delta_{\mathbf{k}, \mathbf{0}}^K. \quad (13)$$

The ensemble average of $\delta^f(\mathbf{k}_1)\delta^{f*}(\mathbf{k}_2)$ then reads:

$$\begin{aligned} \langle \delta^f(\mathbf{k}_1)\delta^{f*}(\mathbf{k}_2) \rangle &= \frac{1}{N^2} \int_{V_\mu} \Pi\left(\frac{\mathbf{r}_1}{H}\right) \Pi\left(\frac{\mathbf{r}_2}{H}\right) \left[\sum_{i \neq j} \langle n_i n_j \rangle W(\mathbf{r}_{i1}) W(\mathbf{r}_{j2}) + \sum_i \langle n_i \rangle W(\mathbf{r}_{i1}) W(\mathbf{r}_{i2}) \right] \\ &\quad \times e^{i\mathbf{r}_1 \cdot \mathbf{k}_1 - i\mathbf{r}_2 \cdot \mathbf{k}_2} d\mathbf{r}_1 d\mathbf{r}_2 - \frac{1}{N} \int_{V_\mu} \Pi\left(\frac{\mathbf{r}_1}{H}\right) \sum_i \langle n_i \rangle W(\mathbf{r}_{i1}) e^{i\mathbf{r}_1 \cdot \mathbf{k}_1} \delta_{\mathbf{k}_2, \mathbf{0}}^K d\mathbf{r}_1 \\ &\quad - \frac{1}{N} \int_{V_\mu} \Pi\left(\frac{\mathbf{r}_2}{H}\right) \sum_i \langle n_i \rangle W(\mathbf{r}_{i2}) e^{-i\mathbf{r}_2 \cdot \mathbf{k}_2} \delta_{\mathbf{k}_1, \mathbf{0}}^K d\mathbf{r}_2 + \delta_{\mathbf{k}_1, \mathbf{0}}^K \delta_{\mathbf{k}_2, \mathbf{0}}^K, \end{aligned} \quad (14)$$

where $\mathbf{r}_{ij} = \mathbf{r}_i - \mathbf{r}_j$. Using

$$\Pi(\mathbf{k}) = \frac{1}{V_\mu} \int_{V_\mu} \Pi\left(\frac{\mathbf{r}}{H}\right) e^{i\mathbf{r} \cdot \mathbf{k}} d\mathbf{r} = \sum_{\mathbf{n}} \delta_{\mathbf{k}, 2k_N \mathbf{n}} \quad (15)$$

where $k_N = \pi/H$ is the Nyquist wavenumber, one can find:

$$\langle \delta^f(\mathbf{k}_1)\delta^{f*}(\mathbf{k}_2) \rangle = \sum_{\mathbf{n}_1, \mathbf{n}_2} \left[|W(\mathbf{k}'_1)|^2 P(\mathbf{k}'_1) \delta_{\mathbf{k}'_1, \mathbf{k}'_2}^K + \frac{1}{N} |W(\mathbf{k}'_1)|^2 \delta_{\mathbf{k}'_1, \mathbf{k}'_2}^K \right], \quad (16)$$

where $\mathbf{k}'_i = \mathbf{k}_i + 2k_N \mathbf{n}_i$ and $W(\mathbf{k})$ is the FT of $W(\mathbf{r})$. For $\mathbf{k}_1 = \mathbf{k}_2 = \mathbf{k}$ we obtain our desired result:

$$\langle |\delta^f(\mathbf{k})|^2 \rangle = \sum_{\mathbf{n}} |W(\mathbf{k} + 2k_N \mathbf{n})|^2 P(\mathbf{k} + 2k_N \mathbf{n}) + \frac{1}{N} \sum_{\mathbf{n}} |W(\mathbf{k} + 2k_N \mathbf{n})|^2, \quad (17)$$

where the summation is over all 3D integer vectors $\mathbf{n}$. The meaning of the above equation is very clear. The density convolution introduces the factor $W^2(\mathbf{k})$ both to the power spectrum and to the shot noise ($1/N$). The finite sampling of the convolved density field results in the ‘alias’ sums (i.e., the sums over $\mathbf{n}$). The alias effect is well-known in Fourier theory, but has not been taken seriously in the power spectrum analysis of large-scale clustering in observational cosmology. Both effects of the convolution and the alias are significant near the Nyquist wavenumber k_N , see Figs. 1 & 2 in the next section.

3. A PROCEDURE TO RECOVER $P(K)$

In the practical measurement of $P(k)$ using FFT, one should first choose a mass assignment function. The NGP ($p = 1$), CIC ($p = 2$) and TSC ($p = 3$) assignment functions

are the most popular function for this purpose. For these schemes, we have (Hockney & Eastwood 1981):

$$W(\mathbf{k}) = \left[\frac{\sin(\frac{\pi k_1}{2k_N}) \sin(\frac{\pi k_2}{2k_N}) \sin(\frac{\pi k_3}{2k_N})}{(\frac{\pi k_1}{2k_N})(\frac{\pi k_2}{2k_N})(\frac{\pi k_3}{2k_N})} \right]^p, \quad (18)$$

where k_i is the i -th component of $\mathbf{k}$.

Once one has selected the assignment function, the shot-noise effect [the second term on the *rhs* of Eq.(17)] is easy to correct. For the NGP, CIC, and TSC assignments, the shot noise term can be expressed by:

$$D^2(\mathbf{k}) \equiv \frac{1}{N} \sum_{\mathbf{n}} W^2(\mathbf{k} + 2k_N \mathbf{n}) = \frac{1}{N} C_1(\mathbf{k}), \quad (19)$$

where $C_1(\mathbf{k})$ are simple analytical functions:

$$C_1(\mathbf{k}) = \begin{cases} 1, & \text{NGP;} \\ \Pi_i[1 - \frac{2}{3} \sin^2(\frac{\pi k_i}{2k_N})], & \text{CIC;} \\ \Pi_i[1 - \sin^2(\frac{\pi k_i}{2k_N}) + \frac{2}{15} \sin^4(\frac{\pi k_i}{2k_N})], & \text{TSC.} \end{cases} \quad (20)$$

Furthermore can easily find out that the $C_1(\mathbf{k})$ of the CIC and TSC schemes are approximately isotropic for $k \leq k_N$, i.e.

$$C_1(\mathbf{k}) \approx \begin{cases} [1 - \frac{2}{3} \sin^2(\frac{\pi k}{2k_N})], & \text{CIC;} \\ [1 - \sin^2(\frac{\pi k}{2k_N}) + \frac{2}{15} \sin^4(\frac{\pi k}{2k_N})], & \text{TSC.} \end{cases} \quad (21)$$

We have tested Equations (19-21) by calculating $\langle |\delta^f(\mathbf{k})|^2 \rangle$ for 10 random simulation samples, each of which consists of $N = 10^5$ points randomly distributed in a unit cube. In this case, $D^2(\mathbf{k}) = \langle |\delta^f(\mathbf{k})|^2 \rangle$. In the upper panel of Figure 1, we show the average values and 1σ errors of $\langle D^2(\mathbf{k})N/C_1(\mathbf{k}) \rangle_d$ estimated from the 10 samples, where (and below) $\langle \dots \rangle_d$ means an average over all directions of $\mathbf{k}$. In this calculation, we have used the NGP, CIC and TSC assignment functions, and have used Eq. (20) for $C_1(\mathbf{k})$. The result for each assignment is shown by one symbol in the figure. From Eq. (19), one expects $\langle D^2(\mathbf{k})N/C_1(\mathbf{k}) \rangle_d = 1$ for all the three assignment functions. Clearly the simulation results agree very well with Eq. (20). In the lower panel, we show the results in a slightly different way, i.e. the averages and 1σ errors of $\langle D^2(\mathbf{k})N \rangle_d$. The solid line is $\langle D^2(\mathbf{k})N \rangle_d = 1$ for the NGP assignment. The dotted and dashed lines are the approximate expressions of Eq. (21) for the CIC and TSC assignments. Again we find a very good agreement between Eq. (21) and the results from the random sample. This means that in most applications, one can use Eq. (21) to correct the shot noise in the FFT measurement of the power spectrum.

After correcting the shot noise, our central problem becomes how to extract $P(k)$ from the first term of Eq. (17). Let us consider the correction factor $C_2(k)$ which is defined as

$$C_2(k) = \frac{\left\langle \sum_{\mathbf{n}} W^2(\mathbf{k} + 2k_N \mathbf{n}) P(\mathbf{k} + 2k_N \mathbf{n}) \right\rangle_d}{P(k)}. \quad (22)$$

Since $W(\mathbf{k})$ is a decreasing function, and $P(k)$ is also a decreasing function on the scales $k \gtrsim k_N$, we expect that the alias contribution of large $|\mathbf{n}|$ to $C_2(k)$ is small. In particular, for $k \ll k_N$, any alias contribution is small and we have $C_2(k) \approx W^2(k) \approx 1$ independent of $P(k)$. For $k \sim k_N$, the alias contribution to $C_2(k)$ becomes important, most of which is owing to the alias of $|\mathbf{k} + 2k_N \mathbf{n}| \sim k$. Therefore the dependence of $C_2(k)$ on $P(k)$ is only the shape of $P(k)$ at $k \sim k_N$, i.e. the local slope α_N of $P(k)$ at $k \sim k_N$, $\alpha_N = [\ln P(k)/\ln k]_{k \sim k_N}$.

Since the local slope α_N is unknown *a priori* in practical measurement of $P(k)$, we propose an iterative method to get the correction factor $C_2(k)$. Suppose that we have measured the power spectrum $P_r(k)$:

$$\begin{aligned} P_r(k) &= \left\langle \langle |\delta^f(\mathbf{k})|^2 \rangle - D^2(\mathbf{k}) \right\rangle_d \\ &= \left\langle \sum_{\mathbf{n}} W^2(\mathbf{k} + 2k_N \mathbf{n}) P(\mathbf{k} + 2k_N \mathbf{n}) \right\rangle_d \end{aligned} \quad (23)$$

for $k \leq k_N$. The local slope α_0 of $P_r(k)$ at $k \sim k_N$ is calculated by a power-law fitting to $P_r(k)$ at $0.5k_N \leq k \leq k_N$. Assuming a power-law form k^{α_0} for $P(k)$ of Eq. (20), we calculate $C_2(k, \alpha_0)$ and get $P_0(k) = P_r(k)/C_2(k, \alpha_0)$. Using the local slope α_0 at $0.5k_N \leq k \leq k_N$ of the $P_0(k)$ just obtained, we calculate $C_2(k, \alpha_0)$ and $P_0(k) = P_r(k)/C_2(k, \alpha_0)$ again. This calculation is repeated until $P_0(k)$ (or α_0) converges to some defined accuracy. The converged $P_0(k)$ is our desired $P(k)$.

Success of the above iteration procedure is shown by Figure 2, where we present our measurement of $P(k)$ for a simulation sample of Jing et al. (1995). The sample is five realizations of P³M simulation of a low-density flat universe with $\Omega_0 = 0.2$, $\Lambda_0 = 0.8$ and $h = 1$. The simulation box size is $128 h^{-1} \text{Mpc}$ and the number of simulation particles is $N = 64^3$. Since the details of the simulation are unimportant for our result here, we will not discuss them anymore. The upper panel of the figure shows the raw power spectra $\langle |\delta^f(\mathbf{k})|^2 \rangle$, which are calculated with the NGP, CIC or TSC mass assignment and with 256^3 or 64^3 grid points. As expected, the raw power spectrum depends on the scheme of the mass assignment: higher order mass assignment gives a smaller $\langle |\delta^f(\mathbf{k})|^2 \rangle$ near the Nyquist frequency. At $k \sim k_N^{64}$, the Nyquist wavenumber of 64^3 grid points, the raw power spectra calculated with 256^3 grid points are expected be very close to the true power spectrum $P(k)$ because neither effect of the shot noise [$D^2(k) \ll P(k)$], the convolution [$W(k) \approx 1$] or

the alias [$C_2(k) \approx 1$] is important. The differences at $k \sim k_N^{64}$ between the power spectra calculated with 64^3 grid points and those with 256^3 grid points, show the importance of the effects discussed in this paper. The lower panel of the figure plots the power spectra $P(k)$ which are corrected following the procedure prescribed previously in this section. In this example, we use Eq. (21) to correct the shot noise, and we require that the slope α_0 converge to the accuracy of $|\Delta\alpha_0| \leq 0.02$ in the iterative method of deconvolving the alias summation. For this accuracy, only fewer than 5 iterations are needed in each calculation. The six power spectra $P(k)$, measured with different mass assignments and with different numbers of grid points, agree so well that their curves overlay each other in the figure. The biggest difference between the 64^3 and 256^3 $P(k)$ is at $k = k_N^{64}$, which is less than 4 percent. The result is very encouraging, and it tells us that using the correction procedure prescribed in this paper, one can obtain the true $P(k)$ for $k \leq k_N$ in the FFT measurement, independent of which mass assignment is used.

In the above derivations and discussions, we have assumed, for simplicity, that the sample is uniformly (within Poisson fluctuation) constructed in a cubic volume. However, the above method is also valid for a sample with selection effects (e.g., an extragalactic catalogue). To show this, let us introduce a selection function $S(\mathbf{r})$ which is defined as the observable rate of the sample at position $\mathbf{r}$. If the underlying density distribution is $n(\mathbf{r})$, the density distribution $n_s(\mathbf{r})$ of the sample is then:

$$n_s(\mathbf{r}) = S(\mathbf{r})n(\mathbf{r}). \quad (24)$$

In this case, we define the following transformation for FFT:

$$\delta_s^f(\mathbf{k}) = \frac{1}{N} \sum_g [n_s^f(\mathbf{r}_g) - \bar{n}S(\mathbf{r}_g)] e^{i\mathbf{r}_g \cdot \mathbf{k}}, \quad (25)$$

where $n_s^f(\mathbf{r}_g)$ is the convolved density of $n_s(\mathbf{r})$ at $\mathbf{r}_g$ [cf. Eq. (9)], and $\bar{n}$ is the mean underlying number density. Following the derivation of Eq. (16), we can easily find:

$$\begin{aligned} \langle \delta_s^f(\mathbf{k}_1) \delta_s^{f*}(\mathbf{k}_2) \rangle &= \sum_{\mathbf{n}_1, \mathbf{n}_2} \left[W(\mathbf{k}'_1) W(-\mathbf{k}'_2) \sum_{\mathbf{k}'_3} S(\mathbf{k}'_1 + \mathbf{k}'_3) P(\mathbf{k}'_3) S(-\mathbf{k}'_2 - \mathbf{k}'_3) \right. \\ &\quad \left. + \frac{1}{N} S(\mathbf{k}'_1 - \mathbf{k}'_2) W(\mathbf{k}'_1) W(-\mathbf{k}'_2) \right], \end{aligned} \quad (26)$$

where $S(\mathbf{k})$ is defined as:

$$S(\mathbf{k}) = \frac{1}{\int S(\mathbf{r}) d\mathbf{r}} \int S(\mathbf{r}) e^{i\mathbf{r} \cdot \mathbf{k}} d\mathbf{r}. \quad (27)$$

Because $S(\mathbf{k})$ peaks at $\mathbf{k} \approx 0$, the coupling of the selection and the power spectrum [the first term of Eq. (26)] is important only at small k . How to treat this coupling is a nontrivial

task and beyond the scope of this paper (but see e.g. Peacock & Nicholson 1991; Jing & Valdarnini 1993). However for larger k where the effects discussed here become significant, we have:

$$\langle |\delta^f(\mathbf{k})|^2 \rangle \approx \sum_{\mathbf{k}_3} S^2(\mathbf{k}_3) \sum_{\mathbf{n}} W^2(\mathbf{k} + 2k_N \mathbf{n}) P(\mathbf{k} + 2k_N \mathbf{n}) + \frac{1}{N} \sum_{\mathbf{n}} W^2(\mathbf{k} + 2k_N \mathbf{n}) \quad (28)$$

The difference between Eq. (26) and Eq. (19) is only the factor $\sum_{\mathbf{k}_3} S^2(\mathbf{k}_3)$. Therefore our procedure for correcting the effects of the mass assignment is still valid for observational samples with selection effects. Equation (28) is derived on the assumption that $S(\mathbf{k})$ is compact in $\mathbf{k}$ space. This assumption is valid for many redshift surveys, e.g., the CfA and IRAS redshift surveys. But is not valid for surveys of irregular boundaries, e.g., pencil beam surveys. In addition, current redshift surveys of galaxies already cover a volume of $(1000 h^{-1} \text{Mpc})^3$. Because one is also interested in the clustering information down to scales of $0.1 h^{-1} \text{Mpc}$, a $(10,000)^3$ gridpoint FFT is required to explore the clustering on all scales, which is still difficult to be realized on modern supercomputers. Other estimators, e.g., the FT of the two-point correlation function (Jing & Börner 2004), should be used to determine $P(k)$ on small scales.

4. SUMMARY

In this paper, we have derived for the first time the formula [Eq. (17)] which relates the raw power spectrum $\langle |\delta^f(k)|^2 \rangle$ estimated with FFT to the true power spectrum $P(k)$. The formula shows clearly how the mass assignment modifies the power spectrum. The convolution of the density field with an assignment function $W(\mathbf{r})$ reshapes the power spectrum (including the shot noise spectrum) by multiplying the factor $|W(\mathbf{k})|^2$; the finite sampling of the convolved density field leads to the alias sum. We have described how to reconstruct $P(k)$ from $\langle |\delta^f(k)|^2 \rangle$. For the NGP, CIC and TSC assignment functions, the shot noise $D^2(\mathbf{k})$ can be expressed by simple analytical functions, therefore the shot noise can be easily corrected. To extract $P(k)$ from the alias sum $\sum_{\mathbf{n}} W^2(\mathbf{k} + 2k_N \mathbf{n}) P(\mathbf{k} + 2k_N \mathbf{n})$, we propose an iterative method. The method has been tested by applying it to an N-body simulation sample. Using different numbers of grid points, we have shown that the method can very successfully recover $P(k)$ for all the NGP, CIC and TSC assignment functions.

The author thanks Bhuvnesh Jain for stimulating discussion on an earlier version of the draft. The work is supported in part by NKBRSF (G19990754) and by NSFC (Nos.10125314, 10373012).

REFERENCES

- Baumgart, D.J. & Fry, J.N. 1991, ApJ, 375, 25
- da Costa, L.N., Vogeley, M.S., Geller, M.J., Huchra, J.P., & Park, C. 1994, ApJ, 437, L1
- Davis, M., Efstathiou, G., Frenk, C.S., & White, S.D.M. 1985, ApJ, 292, 371
- Fisher, K., Davis, M., Strauss, M.A., Yahil, A., & Huchra, J.P. 1993, ApJ, 402, 42
- Gramann, M. & Einasto, J. 1992, MNRAS, 254, 453
- Hockney, R.W. & Eastwood, J.W. 1981, Computer simulations using particles. Mc Graw-Hill
- Jing, Y.P., Mo, H.J., Börner, G. & Fang, L.Z. 1995, MNRAS, (submitted)
- Jing, Y.P. & Valdarnini, R. 1993, ApJ, 406, 6
- Jing, Y. P. & Boerner, G. 2004, ApJ (in press); astro-ph/0406077
- Kaiser, N. 1991, in Texas/ESO-CERN Symposium on Relativistic Astrophysics, eds. J. Barrow et al. (New York: New York Academic Science), 295
- Park, C., Gott, J.R., & Da Costa, L.N. 1992, ApJ, 392, L51
- Peacock, J.A. & Nicholson, D. 1991, MNRAS, 253, 307
- Peacock, J.A. & West, M. 1992, MNRAS, 259, 494
- Peebles, P.J.E. 1980, The large scale structure of the universe. (Princeton: Princeton University Press)
- Scoccimarro, R., Colombi, S., Fry, J. N., Frieman, J. A., Hivon, E., & Melott, A. 1998, ApJ, 496, 586
- Vogeley, M.S., Park, C., Geller, M.J., Huchra, J.P. 1992, ApJ, 391, L5




Fig. 1.— The shot noise $D^2(\mathbf{k})$ estimated from 10 samples of Poisson distributed random points. Each symbol represents the result for each mass assignment, as indicated in the figure. The upper panel shows the result with the function $\langle D^2(\mathbf{k})N/C_3(\mathbf{k}) \rangle_d$, i.e. the $D^2(\mathbf{k})$ scaled to $1/NC_1(\mathbf{k})$. The estimated result agrees quite well with the analytical prediction $\langle D^2(\mathbf{k})N/C_3(\mathbf{k}) \rangle_d = 1$. The lower panel compares the estimated $\langle D^2(\mathbf{k})N \rangle_d$ with our analytical predictions for the NGP (solid line), CIC (dotted line) and TSC (dashed line) assignment functions. For CIC and TSC, we have used the approximate formulae of Eq. (21).




Fig. 2.— The six power spectra which we measured for an N-body simulation sample using three mass assignments (NGP, CIC and TSC) and two grids (64^3 and 256^3 grid points). k_N^{64} is the Nyquist wavenumber of 64^3 grid points. The upper panel shows the raw power spectra $\langle |\delta^f(k)|^2 \rangle$ estimated directly from the FFT (Eq. 8). The lower panel shows the true power spectra $P(k)$ reconstructed from the $\langle |\delta^f(k)|^2 \rangle$ following the procedure described in the text. The six reconstructed $P(k)$ agree so well that their curves overlay each other.